Lactate in Labor

Signal, noise, and a one-hour postpartum reset



Dr. Jennifer Lopez DO, FACOG
JPS Health Network, OBGYN Residency Program
Acclaim Health

Why lactate in obstetric sepsis is worth getting right

Early recognition saves lives; indiscriminate interpretation creates noise.

14.3%

U.S. pregnancy-related deaths due to infection/sepsis, 2017–2019

1 hr

SMFM: empiric broad-spectrum antibiotics ideally within 1 hour for septic shock or high likelihood of sepsis

Trajectory

A repeat value can separate transient exertional physiology from persistent hypoperfusion

Lactate helps identify occult tissue hypoperfusion and risk.
Pregnancy and labor change vital signs, WBC count, respiratory physiology, and metabolic load.
The practical problem: “elevated in labor” ≠ “benign” and ≠ “sepsis” without context.

Sources: Sepsis Alliance/CDC pregnancy-related mortality data; SMFM Consult Series #67; Surviving Sepsis Campaign 2021.

Start with the bedside question

A common labor scenario exposes the interpretive problem.

Laboring patient

39 weeks
2 hours of pushing
T 38.2°C
HR 125 bpm
BP 112/68
Lactate 3.2 mmol/L

3.2

mmol/L

Question 1

Is this evolving sepsis, labor physiology, or both?

Question 2

What additional information would make the number more interpretable?

The talk is about creating that additional information.

The clinical challenge

Labor is not a resting physiology state.

Sepsis signal
Hypoperfusion • mitochondrial stress • impaired clearance

Labor noise
Uterine + skeletal muscle work • catecholamines • dehydration • hemorrhage

Clinical risk
Over-triage burns trust; under-triage misses deterioration

Sepsis physiology

Labor physiology

Bedside context

The same biomarker can represent very different physiology.

Where can lactate come from during labor?

Hyperlactatemia has overlapping physiologic and pathologic drivers.

Labor-related contributors

- Uterine + skeletal muscle work
- Catecholamine-driven glycolysis
- Reduced intake / relative dehydration
- Prolonged pushing or exertion
- Pain and adrenergic stress


Pathologic contributors

- Sepsis and impaired perfusion
- Hemorrhage / hypovolemia
- Hypoxemia or severe anemia
- Seizure activity
- Impaired hepatic clearance

The overlap is why a single intrapartum value can be difficult to classify.

What lactate means in sepsis

A severity marker, not a diagnosis by itself.



Sepsis care bundles treat elevated lactate as a cue to intensify assessment and resuscitation.

Measure lactate when sepsis is suspected or identified in pregnancy/postpartum.
Interpret in clinical context and pursue lactate decrease when elevated.
Do not delay antibiotics, cultures, fluids, vasopressors, or source control while clarifying physiology.

Danger threshold often operationalized at ≥4 mmol/L

Sources: SMFM Consult Series #67; Surviving Sepsis Campaign 2021.

What we know about lactate in labor

Elevations are common; very high values are less common.

40.8%

had lactate ≥2 mmol/L in a labor/puerperium reference study

95.3%

had lactate <4 mmol/L in that same reference cohort

Caution

peripartum lactate is not consistently associated with infection markers in available studies




The interpretive gap is most relevant for lactate 2–3.9 mmol/L.
A value ≥4 mmol/L remains uncommon enough in uncomplicated labor to warrant serious evaluation.
The field lacks pregnancy-specific data on optimal repeat timing and clearance targets.

Sources: Dockree et al., BJOG 2022; Nguyen et al., J Clin Gynecol Obstet 2023; Aguilera et al., J Midwifery Womens Health 2024.

Why a 1-hour postpartum repeat is biologically plausible

Delivery ends a major exertional stimulus. Lactate should move quickly if physiology is transient.



Exercise data: blood lactate can return toward baseline within ~60–80 minutes, depending on recovery conditions.

Labor analogy: uterine/skeletal muscle work stops or markedly falls after birth — but hemorrhage, infection, shock, and hepatic dysfunction do not.


Therefore: the postpartum trend may be more clinically interpretable than the intrapartum value alone.

Sources: Devlin et al., 2014 lactate-clearance exercise physiology; Bester et al., 2023 marathon metabolic recovery.

Exercise analogy is hypothesis-generating; obstetric validation is the goal.

The working hypothesis

The one-hour postpartum repeat converts a single noisy value into a trajectory.



What matters is not only “how high?” but “what happens when labor ends?”

Rapid normalization supports transient exertional physiology when the patient is clinically improving. Failure to clear, recurrent rise, or ≥4 mmol/L after delivery should trigger urgent reassessment. Trend-based interpretation aligns with adult sepsis principles while respecting labor physiology.

Schematic curves — not local outcomes data.

Same starting lactate. Very different trajectories.

A postpartum repeat adds information that the intrapartum value cannot provide alone.

Patient A

In labor: 3.2 mmol/L
+1 hr postpartum: 1.7 mmol/L
HR improving • normotensive
Clinically well


Patient B

In labor: 3.2 mmol/L
+1 hr postpartum: 4.1 mmol/L
Persistent tachycardia
New hypotension

The initial number is identical. The trajectory changes the risk assessment.

Our lactate workflow

Repeat every lactate collected during labor 1 hour after delivery.




The repeat is not a replacement for sepsis treatment. It is an added safety/interpretation checkpoint.

Source framework: SMFM Consult Series #67; local protocol concept supplied by JPS team.

Interpretation algorithm

Escalate from trajectory + bedside status, not lactate alone.



Thresholds should be aligned with your local sepsis policy before go-live.

Operational design

The science only helps if the redraw reliably happens.

Order logic: lactate drawn while patient is admitted to L&D → automatic postpartum redraw prompt.
Handoff logic: delivery team names the repeat lactate during transfer to postpartum/recovery.
Lab logic: timed order due 60 minutes after recorded delivery time.
Escalation logic: persistent elevation pages charge RN/OB clinician/sepsis champion pathway.

Reliability target: “no intrapartum lactate without a postpartum data point.”



How the repeat helps us classify risk

Four clinically useful trajectories.


Physiologic labor signal



Delayed clearance



Sustained hypoperfusion



Recurrent rise



Schematic patterns adapted from sepsis lactate-clearance principles and labor physiology hypothesis.

Data we are collecting

A QI metric should answer “did this change care?” not just “did we draw the lab?”

Domain	Candidate metric
Reliability	% intrapartum lactates with a +1 hr postpartum repeat completed
Signal	Median absolute and relative lactate change after delivery
Safety	ICU transfer, vasopressor use, severe sepsis/septic shock, SMM
Stewardship	Antibiotic escalation/de-escalation linked to lactate trajectory
Equity	Completion and escalation performance by language, race/ethnicity, payer, unit census

Local baseline to verify:
- **~259 sepsis reviews in May 2025–May 2026**
- **~20 maternal sepsis cases/month**
- **~17% severe sepsis among reviewed sepsis cases**

Best next analytic step: separate lactates by timing: antepartum, active labor, immediate postpartum, ≥1 hr postpartum.

Local baseline numbers are draft/contextual — replace with final chart abstraction before presentation.

Expected value for a hospital sepsis program

Better signal, cleaner escalation, stronger multidisciplinary learning.

1. **Differentiate transient exertional lactatemia from persistent hypoperfusion.**
2. **Give nursing and physician teams a concrete postpartum checkpoint.**
3. **Build an obstetric lactate dataset for local thresholds and outcome prediction.**

Bottom line: we are moving from “one abnormal number” to “a reproducible risk trajectory.”

Guardrails: what a 1-hour repeat must never do

A reassuring trajectory is not a substitute for clinical judgment.

Never delay empiric antibiotics or resuscitation for a patient with septic shock or high likelihood of sepsis.
Do not ignore lactate ≥4 mmol/L, rising lactate, hypotension, altered mental status, oliguria, hypoxia, or escalating oxygen/pressor needs.
Account for noninfectious drivers: hemorrhage, seizures, hepatic dysfunction, severe anemia, beta-agonists, metformin, prolonged hypoxia.
If uterine source is suspected, prompt delivery/evacuation is source control — regardless of gestational age.

The repeat lactate is an additional safety net — not permission to watch an unstable patient.

Source: SMFM Consult Series #67; Surviving Sepsis Campaign 2021.

What we do not know yet

The protocol is a testable QI hypothesis—not a validated obstetric cutoff.

Timing

Is +60 minutes the optimal postpartum sampling point?

How much does delivery mode or duration of pushing change clearance?

Threshold

What absolute value or percent decline is truly reassuring?

Should the 2–3.9 mmol/L gray zone be handled differently?

Outcomes

Does trajectory improve escalation, stewardship, ICU prediction, or severe maternal morbidity?

Can we reduce false-positive alarms safely?

The purpose of the QI dataset is to answer these questions rather than assume the answers.

What we need from the broader sepsis program

Standardize the checkpoint; study the signal.

Protocol alignment

Embed +1 hr postpartum lactate repeat into sepsis policy and order sets.

Education

Teach lactate trajectory using labor-specific scenarios.

Dashboard

Track completion, clearance, escalation, and outcomes monthly.

Research/QI

Validate thresholds, clearance targets, and false-positive reduction.

Ask: make postpartum lactate trajectory a routine part of obstetric sepsis review.

Take-home framework

Lactate is a danger signal.

Labor adds physiologic noise.

The +1 hr postpartum repeat adds trajectory.



The goal is earlier escalation for the right patients — and less noise for everyone else.

Selected references

1. Society for Maternal-Fetal Medicine Consult Series #67: Maternal sepsis. Am J Obstet Gynecol. 2023.
2. Evans L, et al. Surviving Sepsis Campaign: International guidelines for management of sepsis and septic shock 2021. Intensive Care Med. 2021.
3. Dockree S, et al. How should we interpret lactate in labour? A reference study. BJOG. 2022.
4. Nguyen M, et al. Association of Labor and Lactate Levels. J Clin Gynecol Obstet. 2023.
5. Aguilera MN, et al. Association of peripartum lactate levels with infection-related outcomes. J Midwifery Womens Health. 2024.
6. Devlin J, et al. Blood lactate clearance after maximal exercise depends on active recovery intensity. 2014.
7. Goyal P, et al. Serial serum lactic acid in pregnancy-associated sepsis for maternal outcome. 2020.
8. Alliance for Innovation on Maternal Health. Sepsis in Obstetric Care Patient Safety Bundle. 2023.